

A REMARK ON ASYMPTOTIC OF HIGHEST WEIGHTS IN TENSOR POWERS OF A REPRESENTATION

KIUMARS KAVEH

ABSTRACT. We consider the semigroup S of highest weights appearing in tensor powers $V^{\otimes k}$ of a finite dimensional representation V of a connected reductive group. We describe the cone generated by S as the cone over the weight polytope of V intersected with the positive Weyl chamber. From this we get a description for the asymptotic of the number of highest weights appearing in $V^{\otimes k}$ in terms of the volume of this polytope.

CONTENTS

1. Introduction	1
2. Semigroups of integral points and convex bodies	3
3. Main result	4
4. Relation with moment polytope	6
References	7

Key words: Reductive group representation, tensor power, highest weight, semigroup of integral points, Newton-Okounkov body, weight polytope, moment polytope, PRV theorem.

AMS subject classification: 05E10, 20G05.

1. INTRODUCTION

Let G be a connected reductive algebraic group over an algebraically closed field $\mathbf{k}$ of characteristic 0, and let V be a finite dimensional G -module. We consider the semigroup of dominant weights

$$S = S(V) = \{(k, \lambda) \mid V_\lambda \text{ appears in } V^{\otimes k}\},$$

where V_λ is the irreducible representation with highest weight λ . In this note we describe the cone $C(S)$ of this semigroup, i.e. the smallest closed convex cone (with apex at the origin) containing S (in other words, the closure of the convex hull of $S \cup \{0\}$). We use this to describe the asymptotic of the number of highest weights λ appearing in $V^{\otimes k}$.

This work is in the spirit of the theory of semigroups of integral points and Newton-Okounkov bodies developed in [Kaveh-Khovanskii09, Part I].

Let A denote the finite set of highest weights in V , i.e. the dominant weight λ for which V_λ appears in V . Consider the union of all the Weyl group orbits of $\lambda \in A$ and let $P^+(V)$ be its convex hull intersected with the positive Weyl chamber. We show that the slice of the cone $C(S)$ at $k = 1$ coincides with the polytope $P^+(V)$ (Theorem 3.1). Our main tool in the proof will be the PRV theorem on the tensor product of irreducible representations.

Let $H_V(k)$ denote the number of dominant weights λ where V_λ appears in $V^{\otimes k}$. From general statements about semigroups of integral points we then conclude that $H_V(k)$ grows of degree $q = \dim(P^+(V))$, i.e. the limit

$$a_q = \lim_{k \rightarrow \infty} H_V(k)/k^q$$

exists and is non-zero. In addition a_q is equal to the (properly normalized) volume of the polytope $P^+(V)$.

In the last section we discuss the connection between the semigroup $S(V)$, its associated polytope $P^+(V)$ with the moment polytope of G -varieties.

At the end, we would like to mention the related paper of Tate and Zelditch [Tate-Zelditch04] in which the authors address the (more difficult) question of describing the asymptotic of multiplicities of irreducible representations appearing in tensor powers of an irreducible representation.

Acknowledgement: the author would like to thank Askold Khovanskii and Kevin Purbhoo for helpful discussions, as well as Shrawan Kumar for helpful email correspondence.

Notation: Throughout the paper we will use the following notation. G denotes a connected reductive algebraic group over an algebraically closed field $\mathbf{k}$ of characteristic 0.

- We fix a Borel subgroup B and a maximal torus T in G . $R = R(G, T)$ is the root system with $R^+ = R^+(G, T)$ the subset of positive roots for the choice of B . The Weyl group of (G, T) is denoted by W . It contains a unique longest element denoted by w_0 .
- Λ denotes the weight lattice of G (that is, the character group of T), and Λ^+ is the subset of dominant weights (for the choice of B). Put $\Lambda_{\mathbb{R}} = \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$. Then the convex cone generated by Λ^+ in $\Lambda_{\mathbb{R}}$ is the positive Weyl chamber $\Lambda_{\mathbb{R}}^+$.
- For a weight $\lambda \in \Lambda$, the irreducible G -module corresponding to λ will be denoted by V_λ and a highest weight vector in V_λ will be denoted by v_λ . Finally for a dominant weight λ , we put $\lambda^* = -w_0(\lambda)$ which is again a dominant weight. One has $V_{\lambda^*} \cong V_\lambda^*$ as G -modules.

2. SEMIGROUPS OF INTEGRAL POINTS AND CONVEX BODIES

Let $S \subset \mathbb{Z}_{\geq 0} \times \mathbb{Z}^n$ be a semigroup of integral points (i.e. S is closed under addition).

Let $C(S)$ be the smallest closed convex cone (with apex at the origin) containing S . Also let $G(S)$ be the subgroup of $\mathbb{Z}^{n+1}$ generated by S and $L(S)$ the linear subspace of $\mathbb{R}^{n+1}$ spanned by S . The sets $C(S)$ and $G(S)$ lie in $L(S)$. To S we associate its *regularization* which is the semigroup $\text{Reg}(S) = C(S) \cap G(S)$. The regularization $\text{Reg}(S)$ is a simpler semigroup with more points which contains the semigroup S . In [Kaveh-Khovanskii09, Section 1.1] it is proved that *the regularization $\text{Reg}(S)$ asymptotically approximates the semigroup S* . More precisely:

Theorem 2.1 (Approximation theorem). *Let $C' \subset C(S)$ be a strongly convex cone which intersects the boundary (in the topology of the linear space $L(S)$) of the cone $C(S)$ only at the origin. Then there exists a constant $N > 0$ (depending on C') such that each point in $G(S) \cap C'$ whose distance from the origin is bigger than N belongs to S .*

Let $\pi : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ denote the projection on the first factor. We call a semigroup $S \subset \mathbb{Z}_{\geq 0} \times \mathbb{Z}^n$ a *non-negative semigroup* if it is not contained in the hyperplane $\pi^{-1}(0)$. If in addition the cone $C(S)$ intersects the hyperplane $\pi^{-1}(0)$ only at the origin, S is called a *strongly non-negative semigroup*.

Let $S_k = S \cap \pi^{-1}(k)$ be the set of points in S at level k . For simplicity throughout this section we assume that $S_1 \neq \emptyset$.

We denote the group $G(S) \cap \pi^{-1}(0)$ by $\Lambda(S)$ and call it the *lattice associated to the non-negative semigroup S* . Finally, the number of points in S_k is denoted by $H_S(k)$. H_S is called the *Hilbert function of the semigroup S* .

Definition 2.2 (Newton-Okounkov convex set). We call the projection of the convex set $C(S) \cap \pi^{-1}(1)$ to $\mathbb{R}^n$ (under the projection on the second factor $(1, x) \mapsto x$), the *Newton-Okounkov convex set of the semigroup S* and denote it by $\Delta(S)$. In other words,

$$\Delta(S) = \overline{\text{conv}\left(\bigcup_{k>0} \{x/k \mid (k, x) \in S_k\}\right)}.$$

If S is strongly non-negative then $\Delta(S)$ is compact and hence a convex body.

Let $\Lambda \subset \mathbb{R}^n$ be a lattice of full rank n . Let $E \subset \mathbb{R}^n$ be a subspace of dimension q which is rational with respect to Λ . The *Lebesgue measure normalized with respect to the lattice Λ* in E is the Lebesgue measure $d\gamma$ in E normalized such that the smallest measure of a q -dimensional parallelepiped with vertices in $E \cap \Lambda$ is equal to 1. The measure of a subset $A \subset E$ will be called its *normalized volume* and denoted by $\text{Vol}_q(A)$ (whenever the lattice Λ is clear from the context).

Let H_S and $H_{\text{Reg}(S)}$ be the Hilbert functions of S and its regularization respectively. From Theorem 2.1 it follows that $H_S(k)$ and $H_{\text{Reg}(S)}(k)$ have the same asymptotic as k goes to infinity. Thus the Newton-Okounkov

convex set $\Delta(S)$ is responsible for the asymptotic of the Hilbert function of S ([Kaveh-Khovanskii09, Section 1.4]):

Theorem 2.3. *The function $H_S(k)$ grows like $a_q k^q$ where q is the dimension of the convex body $\Delta(S)$. This means that the limit*

$$a_q = \lim_{k \rightarrow \infty} H_S(k)/k^q$$

exists and is non-zero. Moreover, the q -th growth coefficient a_q is equal to $\text{Vol}_q(\Delta(S))$, where the volume is normalized with respect to the lattice $\Lambda(S)$.

Finally we prove a proposition which will be used later in proof of the main result (Theorem 3.1).

Proposition 2.4. *Let $S \subset \mathbb{Z}_{\geq 0} \times \mathbb{Z}^n$ be a non-negative semigroup and $C = C(S)$ the cone associated to S . Let $C' \subset C$ be a convex cone of full dimension (centered at the origin) and $S' = S \cap C'$ the subsemigroup consisting of all the points of S contained in C' . Then the cone $C(S')$ associated to S' coincides with C' .*

Proof. Clearly $C(S') \subset C'$. By contradiction suppose $C(S')$ is not equal to C' . Then there is a convex cone $\tilde{C} \subset C'$ of full dimension which does not intersect $C(S')$ and moreover intersects the boundary of C' (in the topology of the subspace $L(S)$) only at the origin. Since $\tilde{C}$ has full dimension it contains a rational point (with respect to the lattice $\Lambda(S)$) which then implies that it contains a point in $\Lambda(S)$. Now applying Theorem 2.1 we see that the convex cone $\tilde{C}$ should contain a point in S' which contradicts that $C(S') \cap \tilde{C} = \emptyset$. $\square$

In the rest of the paper we will deal with semigroups and convex polytopes naturally associated to a reductive group G and its representations.

3. MAIN RESULT

Let V be a finite dimensional G -module. Define the set $S(V) \subset \mathbb{Z}_{\geq 0} \times \Lambda^+$ by

$$S(V) = \{(k, \lambda) \mid V_\lambda \text{ appears in } V^{\otimes k}\}.$$

If v_λ and v_μ are highest weight vectors in $V^{\otimes k}$ and $V^{\otimes \ell}$ with weights λ and μ respectively, then $v_\lambda \otimes v_\mu$ is a highest weight vector in $V^{\otimes k+\ell}$ of weight $\lambda + \mu$. It follows that $S(V)$ is a semigroup with respect to addition. Let $\Delta(V)$ denote the Newton-Okounkov body of the semigroup $S(V)$. In other words,

$$\Delta(V) = \overline{\text{conv}\left(\bigcup_{k>0} \{\lambda/k \mid V_\lambda \text{ appears in } V^{\otimes k}\}\right)}.$$

Also let A be the collection of γ where V_γ appears in V . Then the *weight polytope* of V is defined by $P(V) = \text{conv}\{w(\gamma) \mid w \in W, \gamma \in A\}$, i.e. the convex hull of the union of Weyl orbits of $\gamma \in A$. We will denote the intersection of $P(V)$ with the positive Weyl chamber $\Lambda_{\mathbb{R}}^+$ by $P^+(V)$ and call it the *positive weight polytope*.

Theorem 3.1 (Main result). $\Delta(V)$ coincides with the positive weight polytope $P^+(V)$.

We will need the following well-known fact:

Lemma 3.2. *Let λ_1, λ_2 be dominant weights and let V_γ appear in $V_{\lambda_1} \otimes V_{\lambda_2}$. Then $\gamma = \lambda_1 + \lambda_2 - \sum_{\alpha \in R^+} c_\alpha \alpha$ where $c_\alpha \geq 0$. From this it follows that γ belongs to $P^+(\lambda_1 + \lambda_2)$.*

Our main tool to prove Theorem 3.1 will be the well-known PRV theorem regarding the tensor product of two irreducible representations. It was conjectured by K. Parthasarathy, R. Ranga Rao and V. Varadarajan in [PRV67]. Later it was proved by S. Kumar in [Kumar88]. We briefly recall its statement.

Theorem 3.3 (PRV theorem). *Let $\lambda_1, \lambda_2 \in \Lambda^+$ be two dominant weights. Suppose for two Weyl group elements $w_1, w_2 \in W$ we have $\gamma = w_1(\lambda_1) + w_2(\lambda_2)$ is a dominant weight. Then V_γ appears in the decomposition of the tensor product $V_{\lambda_1} \otimes V_{\lambda_2}$ into irreducible representations.*

Define the set $\tilde{S}(V) \subset \mathbb{Z}_{\geq 0} \times \Lambda$ by:

$$\tilde{S}(V) = \{(k, w(\lambda)) \mid w \in W, V_\lambda \text{ appear in } V^{\otimes k}\}.$$

Roughly, speaking $\tilde{S}(V)$ is the union of W -otbits of λ for which V_λ appears in some tensor power $V^{\otimes k}$. Notice that $S(V) = \tilde{S}(V) \cap (\mathbb{Z}_{\geq 0} \times \Lambda^+)$. The following is a straight forward corollary of Theorem 3.3

Corollary 3.4. *1) $\tilde{S}(V)$ is a semigroup. 2) The convex body $\Delta(\tilde{S}(V))$ associated to this semigroup coincides with $P(V)$.*

Proof. 1) Let $(k, w_1(\lambda_1)), (\ell, w_2(\lambda_2))$ be two elements in $\tilde{S}(V)$. We can write $w_1(\lambda_1) + w_2(\lambda_2)$ as $w(\lambda)$ for some $\lambda \in \Lambda, w \in W$. By Theorem 3.3, V_λ appears in $V_{\lambda_1} \otimes V_{\lambda_2}$ and hence it appears in $V^{\otimes \ell+k}$. This shows that $(\ell + k, w_1(\lambda_1) + w_2(\lambda_2)) = (\ell + k, w(\lambda))$ belongs to $\tilde{S}(V)$ which proves 1). 2) By Lemma 3.2, for any integer $k > 0$, the convex hull of $\tilde{S}(V)_k$ is $kP(V)$ and hence $\Delta(\tilde{S}(V))$ coincides with $P(V)$. $\square$

Proof of Theorem 3.1. From Proposition 2.4, the convex body $\Delta(V)$ associated to the semigroup $S(V) \subset \tilde{S}(V)$ is just the intersection of $\Delta(\tilde{S}(V))$ with $\Lambda_{\mathbb{R}}^+$. By Corollary 3.4(2) we know that $\Delta(\tilde{S}(V))$ is the weight polytope $P(V)$ which finishes the proof. $\square$

Corollary 3.5. *Let $H_V(k)$ be the number of λ such that V_λ appears in $V^{\otimes k}$. Then $H_V(k)$ grows of degree $q = \dim P^+(V)$. That is, the limit*

$$a_q = \lim_{k \rightarrow \infty} H_V(k)/k^q$$

exists and is non-zero. Moreover, a_q is equal to $\text{Vol}_q(P^+(V))$, where volume is the Lebesgue measure in $\Lambda_{\mathbb{R}}$ normalized with respect to the lattice $\Lambda(S(V)) \subset \Lambda$.

Proof. Follows directly from Theorem 3.1 and Theorem 2.3. $\square$

4. RELATION WITH MOMENT POLYTOPE

In this section we see how the positive weight polytope $P^+(V)$ appears as a moment polytope for the action of $G \times G$ on G .

Let V be a finite dimensional G -module, and $X \subset \mathbb{P}(V)$ an irreducible closed G -invariant subvariety. Let $R = \bigoplus_{k \geq 0} R_k$ denote the homogeneous coordinate ring of X . It is a graded G -algebra. Following Brion, one defines the *moment polytope* $\Delta(X)$ to be:

$$\Delta(X) = \overline{\text{conv}\left(\bigcup_{k > 0} \{\lambda/k \mid V_\lambda \text{ appears in } R_k\}\right)}.$$

One shows that $\Delta(X) \subset \Lambda_{\mathbb{R}}^+$ is a polytope (see [Brion87]). Moreover, when $\mathbf{k} = \mathbb{C}$ and X is smooth, the polytope $\Delta(X)$ can be identified with the moment polytope of X regarded as a Hamiltonian space for the action of a maximal compact subgroup K of G and the symplectic structure induced from the projective space (see for example [Guillemin-Sternberg84]).

Let $G \times G$ act on G via multiplication from left and right. Let $\mathbf{k}[G]$ denote the algebra of regular functions on the variety G . It is a rational $G \times G$ -module. It is well-known that for each dominant weight λ , the (λ^*, λ) -isotypic component $\mathbf{k}[G]_{(\lambda^*, \lambda)}$ is isomorphic to $V_{\lambda^*} \otimes V_\lambda$. Moreover, any isotypic component of $\mathbf{k}[G]$ is of this form for some λ . In fact any $G \times G$ -isotypic component $\mathbf{k}[G]_{(\lambda^*, \lambda)}$ in $\mathbf{k}[G]$ is the linear span of the matrix entries corresponding to the representation of G in V_λ (see [Kraft85]). Now if $\lambda_1, \lambda_2 \in \Lambda^+$ are two dominant weights, the product $\mathbf{k}[G]_{(\lambda_1^*, \lambda_1)} \mathbf{k}[G]_{(\lambda_2^*, \lambda_2)}$ is the linear span of the matrix entries corresponding to $V_{\lambda_1} \otimes V_{\lambda_2}$. This shows that we have the following decomposition for the product of isotypic components in the algebra $\mathbf{k}[G]$:

$$(1) \quad \mathbf{k}[G]_{(\lambda_1^*, \lambda_1)} \mathbf{k}[G]_{(\lambda_2^*, \lambda_2)} = \bigoplus_{\gamma \in \chi(\lambda_1, \lambda_2)} V_{\gamma^*} \otimes V_\gamma,$$

where $\chi(\lambda_1, \lambda_2)$ denotes the collection of all $\gamma \in \Lambda^+$ for which V_γ appears in $V_{\lambda_1} \otimes V_{\lambda_2}$.

Now let $\pi : G \rightarrow \text{GL}(V)$ be a finite dimensional representation. Then $\text{End}(V)$ is naturally a $G \times G$ -module where $G \times G$ acts via π by multiplication from left and right. Let $\tilde{\pi} : G \rightarrow \mathbb{P}(\text{End}(V))$ be the induced map to projective space and let X be the closure of the image of G in $\mathbb{P}(\text{End}(V))$. It is a $G \times G$ -invariant closed irreducible subvariety.

From (1) one can see the following:

Proposition 4.1. *Let $R = \bigoplus_k R_k$ denote the homogeneous coordinate ring of X in $\mathbb{P}(\text{End}(V))$. Then for $k > 0$ we have: $V_{(\lambda^*, \lambda)}$ appears in R_k if and only if V_λ appears in $V^{\otimes k}$. It follows that, under the projection on the second factor, $\Delta(X) \subset \Lambda_{\mathbb{R}}^+ \times \Lambda_{\mathbb{R}}^+$ identifies with $P^+(V)$.*

Remark 4.2. The relation between the moment polytope of X (i.e. a group compactification) and the weight polytope $P^+(V)$ has also been shown in [Kazarnovskii87] using methods from symplectic geometry.

REFERENCES

- [Brion87] Brion, M. *Sur l'image de l'application moment*. Sminaire d'algbre Paul Dubreil et Marie-Paule Malliavin (Paris, 1986), 177–192, Lecture Notes in Math., 1296, Springer, Berlin, 1987.
- [Guillemin-Sternberg84] Guillemin, V.; Sternberg, S. *Geometric quantization and multiplicities of group representations*. Invent. Math. 77 (1984), 533–546.
- [Kaveh-Khovanskii09] Kaveh, K.; Khovanskii, A. G. *Newton-Okounkov convex bodies, semigroups of integral points, graded algebras and intersection theory*. Preprint: arXiv:0904.3350v1
- [Kazarnovskii87] Kazarnovskii, B. *Newton polyhedra and the Bezout formula for matrix-valued functions of finite dimensional representations*. Functional Analysis and its applications, v. 21, no. 4, 73–74 (1987).
- [Kraft85] Kraft, H. *Geometrische Methoden in der Invariantentheorie*. (Geometrical methods in invariant theory). Aspects of Mathematics, D1. Friedr. Vieweg & Sohn, Braunschweig, 1984.
- [Kumar88] Kumar, S. *Proof of the Parthasarathy-Ranga Rao-Varadarajan conjecture*. Invent. Math. 93 (1988), no. 1, 117–130.
- [PRV67] Parthasarathy, K. R.; Ranga Rao, R.; Varadarajan, V. S. *Representations of complex semi-simple Lie groups and Lie algebras*. Ann. of Math. (2) 85 (1967) 383–429.
- [Tate-Zelditch04] Tate, T.; Zelditch, S. *Lattice path combinatorics and asymptotics of multiplicities of weights in tensor powers*. J. Funct. Anal. 217 (2004), no. 2, 402–447.